



Civilian Employment After Military Service

Jaime L. Horton

Isabel G. Jacobson

Edward J. Boyko

Charlene A. Wong

Besa Smith

Timothy S. Wells

Margaret A.K. Ryan

Tyler C. Smith

for the Millennium Cohort Study Team



Naval Health Research Center

Report No. 11-49

The views expressed in this article are those of the authors and do not necessarily reflect the official policy or position of the Department of the Navy, Department of Defense, nor the U.S. Government.

Approved for public release; distribution unlimited.

This research was conducted in compliance with all applicable federal regulations governing the protection of human subjects in research.

***Naval Health Research Center
140 Sylvester Road
San Diego, California 92106-3521***

ORIGINAL ARTICLE

The impact of prior deployment experience on civilian employment after military service

Jaime L Horton,¹ Isabel G Jacobson,¹ Charlene A Wong,¹ Timothy S Wells,¹ Edward J Boyko,² Besa Smith,¹ Margaret A K Ryan,³ Tyler C Smith,¹ for the Millennium Cohort Study Team

¹Deployment Health Research Department, Naval Health Research Center, San Diego, California, USA

²Department of Veterans Affairs Puget Sound Health Care System, Seattle Epidemiologic Research and Information Center, Seattle, Washington, USA

³Clinical Investigation Program, Naval Hospital Camp Pendleton, Camp Pendleton, California, USA

Correspondence to

Jaime L Horton,
Deployment Health Research Department, Naval Health Research Center, 140 Sylvester Road, San Diego, CA 92106-5122, USA;
Jaime.Horton@med.navy.mil

Received 19 July 2012

Revised 18 December 2012

Accepted 28 February 2013

Published Online First

21 March 2013

ABSTRACT

Objective To determine if deployment to recent military operations or other health, demographic, or military-related characteristics were associated with employment after military service.

Methods Former US active duty military service members participating in the Millennium Cohort Study, a population-based sample of US military personnel that began in July of 2001, were prospectively followed from the time of baseline health reporting to self-reported employment status after military separation.

Results Of the 9099 separated personnel meeting inclusion criteria, 17% reported unemployment after military service. In multivariable modelling, prior deployment experiences, with or without reported combat, and post-traumatic stress disorder (PTSD) were not significantly associated with employment status postservice. Among those who routinely retired from service with a pension, positive screens for depression (OR, 1.67; 95% CI, 1.05 to 2.63) and panic/anxiety (OR, 1.63; 95% CI, 1.10 to 2.43) were significantly associated with subsequent unemployment. Poor physical health, female sex, black race, lower education and disabling illnesses/injuries were also predictive of postservice unemployment.

Conclusions After stratifying for reason for military separation, mental disorders like depression or panic/anxiety and poor physical health may have greater impact than prior deployment experiences or PTSD on the ability to find or maintain employment postservice. These findings may guide support for veterans most in need of job placement services after military service.

Service in the US armed forces offers training, resources for educational advancement, and valuable experience often sought by civilian employers. While service members share common workplace exposures with the general population, they also have unique experiences such as military deployment. Recent US military service, in particular, may be characterised by multiple and prolonged deployments. Service members who deployed during times of war and combat may face obstacles or discrimination when seeking employment in the civilian world due to common misperceptions, associated stereotypes, or readjustment issues.^{1 2}

Stressful experiences while deployed in support of the operations in Iraq and Afghanistan have been shown to increase risk for postdeployment

What this paper adds

- ▶ US service members have many unique experiences, such as multiple and prolonged military deployments, which have been shown to increase risk for postdeployment mental health morbidity.
- ▶ There is growing concern that deployments during the recent conflicts, and the accompanying societal stigma of adverse military experiences, may serve as a barrier to postservice employment.
- ▶ In multivariable modelling, prior deployment experiences and post-traumatic stress disorder (PTSD) were not significantly associated with job status postservice, while screening positive for depression or panic/anxiety or reporting poor physical health was associated with increased risk of postservice unemployment among certain groups.
- ▶ These findings provide a better understanding of the circumstances associated with unemployment postservice among US veterans which may have long-term health and economic impacts.

mental health morbidity that may have a significant, lasting impact on health.^{3–6} A high proportion of services by the department of Veterans Affairs (VA) are for the care of mental disorders; in fact, in 2008, five of the top 20 most frequent discharge diagnoses were for mental disorders or substance abuse.⁷ The burden of care on the VA system may be augmented due to unemployment associated with mental disorders, since lack of employment means no job-provided health insurance and individuals with mental illness often struggle in their efforts to obtain or maintain employment. Thus, there is growing concern that the recent conflicts and the reported increase in mental illness and their accompanying societal stigma may serve as a barrier to postservice employment.^{8 9}

The Millennium Cohort Study was designed to longitudinally examine any impact military service may have on the health and well-being of service members during and after their military careers.^{10–13} Few studies have examined the employability of returning OEF/OIF veterans,^{14 15}

To cite: Horton JL, Jacobson IG, Wong CA, et al. *Occup Environ Med* 2013;**70**:408–417.

and to our knowledge, no studies have examined the association of recent deployment experience prospectively with subsequent civilian employment among former active duty military personnel. The present study prospectively assesses whether civilian employment status differs among cohort members with deployment experience compared with those who did not deploy, and whether the effect of military deployments on employment is modified by mental health. Further, in accordance with VA concerns,¹⁴ these analyses aimed to identify subgroups of the military who report lower rates of employment after separation from service, and who may potentially benefit from additional training, treatment, or support to successfully transition to civilian employment.

METHODS

Study population

The Millennium Cohort Study began collection of self-reported outcome and exposure data in 2001, prior to the start of recent combat operations in Iraq and Afghanistan, and resurveys its participants every 3 years through 2022. Using a phased enrolment strategy, the Millennium Cohort currently includes more than 150 000 members. Invited subjects to the first panel in 2001 included a probability-based sample of 256 400 regular active duty, Reserve, and National Guard members randomly drawn from electronic rosters at the Defense Manpower Data Center (DMDC) as of 1 October 2000, and contacted via postal mail, e-mail and questionnaire mailing to complete a paper or web survey. In 2004, a second panel of 150 000 regular active duty, Reserve and National Guard personnel with 1–2 years of military service as of 1 October 2003 was invited to complete a survey on the web or by paper. Other modes of contact and tailored messages were used during and between survey cycles, based on a modified Dillman approach,¹⁶ to keep participants engaged in the study. Detailed descriptions of the methodology for the Millennium Cohort Study have been published elsewhere.^{10–12}

The first panel of 77 047 consenting participants (36.0% of those believed to have been contacted) enrolled in 2001–2003. The first follow-up questionnaire (2004–2006) was completed by 55 021 members (71% response rate), and the second follow-up (2007–2008) was completed by 54 790 members (71%). The second panel (2004–2006) enrolled 31 110 consenting participants (25.4% of those believed to have been contacted), and 17 152 members (55%) completed one follow-up questionnaire (2007–2008). To be included in the current study, panels 1 and 2 participants must have been serving in the military on active duty status at the time of their baseline survey, completed at least one follow-up, and separated from the military between baseline and follow-up. Panel 1 participants who completed two follow-up surveys, had their demographic and health information assessed at the closest survey *prior* to their separation date, while their employment status was determined using the closest survey *after* their separation date.

Separation from the US military

Military separation status was determined using DMDC data, and confirmed with self-report where possible. Nearly all panel 2 participants (99%) were confirmed separated with self-report, compared with 53% of panel 1 participants confirmed with self-report. The question on the survey 'Are you currently serving in the military (active duty, Reserve, or National Guard)?' was not introduced until the 2007–2008 survey period; therefore, panel 1 subjects who separated prior to the first follow-up in 2004–2006 could not be verified with self-report. Among those

with both records, fewer than 3% had discordant results and were excluded to reduce misclassification. DMDC maintains an archive of personnel data covering all service branches, components and phases of the military cycle (accession, separation, retirement). Electronic separation files included date of separation as well as codes that indicate the reason for separation. Participants were divided into three categories based on these codes: (1) those who fulfilled at least 20 years of service and had a pension (routinely retired), (2) those who fulfilled their contract term of service (typically 4 years) and (3) those who separated for all other reasons (eg, disability, parenthood, misconduct, early release). Contractual term of service at enlistment may vary by service branch and component, and may range between two and 6 years.¹⁷ Of note, subjects in this study did not include any service members who separated prior to initial entry training because early separation of this nature did not fulfil inclusion criteria in the Millennium Cohort Study.

Civilian employment

Civilian employment was assessed at the closest survey following the service member's separation date. Employment status was determined by the question in the Millennium Cohort questionnaire, 'Do you have a civilian job at this time?', with the response options of 'Yes', 'No civilian employment at this time,' or 'Homemaker.' Those who reported 'yes' were defined as 'employed' and those who reported 'no civilian employment at this time' were termed 'unemployed' for the purposes of this paper. We could not determine whether these 'unemployed' individuals were actively seeking employment, however, so we could not equate this self-reported employment status to national unemployment rates which only include job seekers. Thus, unemployment rates are not presented for this study population. Those who answered 'Homemaker' were excluded from analyses because the numbers in this group were insufficient for multivariable modelling, and they represented a unique group (younger, married, 89% women) who could not easily be included with the employed or unemployed individuals.

Deployment experience

Military deployment data from DMDC were used to categorise individuals as not recently deployed and deployed with or without combat experience. Those with no deployments prior to military separation or only prior to baseline were considered not recently deployed. Deployed participants must have completed their first deployment between baseline and their date of separation. Combat experience was determined by at least one affirmative response to witnessing the following in the 3 years prior to outcome assessment: a person's death due to war, instances of physical abuse, dead and/or decomposing bodies, maimed soldiers or civilians, or prisoners of war or refugees.

Covariates

Covariates were assessed using the closest survey prior to separation from military service, and included the following: sex, age at separation, education, race/ethnicity, marital status, military pay grade (a surrogate for socioeconomic class), service branch, and military occupation. Time separated was also calculated by the number of days from the date of separation to the follow-up questionnaire. Period of study enrollment (panel 1 or 2) was also included since panel 1 members are proportionally older with more education and a higher pay grade than those in panel 2; attributes that may position them better for employment. Finally, we examined smoking status, alcohol-related problems (≥ 1 problem indicated a positive screen) as defined by the

Patient Health Questionnaire (PHQ),^{18 19} disabling illness or injury (self-reported), life stressors (defined using the Holmes and Rahe Scale²⁰), and physical component summary (PCS) score (assessed using the Short Form 36-Item Health Survey for Veterans (SF-36V)²¹). The SF-36V includes questions about general health, bodily pain and ability to work and carry out daily activities. Participants' health was categorised by percentile of the PCS score based on overall scores of panels 1 and 2 of the Millennium Cohort.

Mental disorders were also assessed using the closest survey prior to separation from military service, and were identified by screening tools embedded in the Millennium Cohort questionnaire. A positive screen for post-traumatic stress disorder (PTSD) was measured using the 17-item PTSD Checklist-Civilian Version.²² Participants screened positive if they reported a moderate or higher level of at least one intrusion symptom, three avoidance symptoms, and two hyperarousal symptoms (criteria established by the *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV-TR)*²³). This approach has been shown to have high sensitivity and high internal consistency in this cohort.^{24 25} A positive screen for depression was assessed using nine items from the PRIME-MD PHQ,¹⁸ corresponding to the depression diagnosis from the *DSM-IV-TR*^{23 26} with sensitivity of 0.93 and specificity of 0.89.²⁷ Screens for panic or other anxiety disorders also used the PHQ for a psychosocial assessment based on scores of several health concepts.^{18 19}

Statistical analyses

Univariate analyses were conducted to evaluate the unadjusted associations between civilian employment and all covariates. Multivariable logistic regression models were built to calculate the odds of unemployment following military service in relation to deployment experience while adjusting for all covariates mentioned. Given the exploratory nature of this study, all defined variables were included. Model diagnostic tests were conducted and we assessed model fit using the Hosmer–Lemeshow goodness-of-fit test. To identify the existence of collinearity, we examined all variance inflation factors and considered a value of four or greater to indicate a potential problem.²⁸ First-order interactions between reason for separation and age at separation, and reason for separation and service branch were tested. Both interactions were significant at $p < 0.10$. Three models were built to examine characteristics of those who left the military for routine retirement, fulfillment of service term, and for all other reasons. To assess whether the association between deployment and post service employment was modified by service members' mental health, we examined the interactions between deployment and each mental health variable in each stratified model. None of these interaction terms was significant ($p < 0.10$); thus, we proceeded with constructing three models not stratified by deployment or mental health variables to estimate the association between prior deployment and postservice employment, adjusting for mental health and all other covariates. Data management and statistical analyses were performed using SAS software, V 9.3 (SAS Institute, Cary, North Carolina, USA).

RESULTS

Of the 80 524 Millennium Cohort members with at least one follow-up assessment, 13 832 separated between their baseline and last submitted follow-up. Participants who selected the 'Homemaker' option ($n=588$) and those missing outcome ($n=450$) or covariate ($n=1720$) information were excluded. Reservists and National Guard members ($n=1975$) were also

removed because many already hold civilian occupations prior to military separation, leaving 9099 participants available for analyses (figure 1).

Characteristics of the population by employment status are shown in table 1. Overall, 17% did not have civilian employment postservice. The majority of the population (60.3%) retired with at least 20 years of service and a pension, 23.8% separated after fulfilling their service term, and 15.9% for other reasons. Mean age at separation was 37.3 years (sd, 9.1 years). χ^2 Tests revealed that positive screens for mental disorders were significantly higher among the unemployed, and a small but significant difference was found across employment categories among those with recent deployment experience. Unemployment was lowest among those who routinely retired (779/5488=14.2%), followed by those who fulfilled their service term (417/2165=19.3%) and those separated for other reasons (383/1446=26.5%).

In adjusted analyses, all covariates were included in the models (unless indicated N/A), and no issues with multicollinearity were detected using the previously described variance inflation factor criterion. All models fit the data well, as indicated by the Hosmer–Lemeshow goodness-of-fit test ($p > 0.10$ for all). Prior deployment experience and PTSD were not associated with post-service unemployment (table 2) in multivariable analyses. Screening positive for depression (OR, 1.67; 95% CI, 1.05 to 2.63) or panic/anxiety (OR, 1.63; 95% CI, 1.10 to 2.43) was associated with significantly higher odds of unemployment, but only among those who routinely retired. Also among the routinely retired, unemployment was significantly associated with being female, reporting a disabling illness or injury, older age at separation, being non-Hispanic black, former Navy or Air Force members, current smokers and reporting poor health. Conversely, current and formerly married, more educated individuals and those separated longer than a year had reduced odds of unemployment.

Among those who separated after fulfilling their service term, reporting a disability, experiencing moderate stress, belonging to the younger cohort (panel 2), being non-Hispanic black, and female, significantly increased the odds of unemployment. Married respondents, former electronic equipment repairers, communications/intelligence specialists, and those separated for longer periods were less likely to be unemployed postservice (table 2).

Those who separated for other reasons were the most likely to be unemployed. Among this group, 48.8% of the unemployed left military service due to a disability, 12.5% left for a pregnancy or parenthood, and most of the remaining members underwent administrative discharge (eg, due to drug abuse, misconduct, unfitness) (data not shown). Characteristics associated with unemployment among those separated for other reasons included older age, being non-Hispanic black, and reporting poor physical health. Unemployment was less likely among those with a bachelor's degree or higher and those separated for longer than a year (table 2).

DISCUSSION

The intense and prolonged deployments during the recent operations in Iraq and Afghanistan, and increases in mental disorders among personnel exposed to combat,^{4 6 29} have given rise to concern over the employability of veterans. To our knowledge, this is the first population-based study to prospectively quantify civilian employment in a large population of former active duty military personnel from all service branches. We found no association between deployment experience and employment status following separation. Routinely retired




Figure 1 Flowchart of Millennium Cohort participants included in the study.

subjects screening positive for depression or panic/anxiety disorders were significantly more likely to report unemployment compared with subjects with no mental disorders. Other characteristics found to be associated with unemployment included black race, female sex, less education and poor physical health.

In this study, deployment experience did not increase risk of unemployment. This finding provides reassurance in light of frequent reference to high unemployment among veterans returning from Iraq and Afghanistan,³⁰ and is consistent with a March 2011 Bureau of Labour Statistics (BLS) news release stating that these veterans had unemployment rates that were not statistically different from veterans who had served elsewhere, nor did veterans who had entered military service since September 2001 have higher unemployment rates than non-veterans of the same age and gender group.³¹ Military members selected for deployment are overall healthier than members who do not deploy;^{3 4 6} thus, there may be other factors, like mental disorders, that are more influential on veterans' ability to obtain civilian employment.

Other important findings included the associations identified between some mental disorders and employment. Depression and anxiety disorders have been associated with work status and productivity in prior studies.^{32 33} A challenge to define cause-effect relationships remains, since mood disorders may predispose one to employment issues, as well as result from unemployment or economic stressors.³⁴ An advantage of the prospective nature of this study was the ability to define mental disorders while participants were still serving in the military, such that these could be determined as factors potentially associated with subsequent unemployment. Our study did not find a significant association between PTSD and subsequent

unemployment, which contrasts with one prior VA study of veterans in a vocational rehabilitation programme,³⁵ a discrepancy likely due to differences in study populations, design and methods. The broad representation of the military in the Millennium Cohort suggests that screening positive for PTSD prior to separation is not a major factor in employability post-service. This finding may be due to the strong support the military and VA have given to members with this 'signature injury,' or it may simply reflect differences in functional impairment between PTSD and mood disorders.³⁶ Also, it should be noted that the association between screening positive for PTSD and employment status existed at the crude level. However, after adjusting for variables such as depression, panic disorder, and physical component score, the association between PTSD and employment status was diminished. This may indicate potential overlap between comorbid mental disorders and physical health impairment that likely play a role, when taken together, on an individual's ability to find or maintain a job.

Additionally, across all groups we noted that those separated for longer time periods were less likely to be unemployed, likely because they had more time to search for a job. Since time was strongly predictive of postservice employment status, we also examined whether time modified the effects of PTSD on employment status. Adding this first-order interaction term into our three multivariable models, we found that it was significant only among those separated for other reasons ($P=0.02$) (data not shown). The significance of the interaction was explained by an association between PTSD and employment status among those separated for more than 2 years whereas no association was found among those separated for 2 years or less. In those separated for more than 2 years, the proportion unemployed

Workplace

Table 1 Characteristics of Millennium Cohort participants who separated from the US military, by civilian employment status

| Baseline* characteristics | Total separated N=9099 (100%) n (%)† | | Unemployed n=1579 (17%) n (%) | | Employed n=7520 (83%) n (%) | | p Value |
|---|---|--------|--|--------|--------------------------------------|--------|---------|
| Reason for separation | | | | | | | |
| Routinely retired | 5488 | (60.3) | 779 | (49.3) | 4709 | (62.6) | <0.001 |
| Fulfilled service term | 2165 | (23.8) | 417 | (26.4) | 1748 | (23.2) | |
| Other reasons | 1446 | (15.9) | 383 | (24.3) | 1063 | (14.1) | |
| Deployment experience | | | | | | | |
| Not recently deployed | 6711 | (73.8) | 1160 | (73.5) | 5551 | (73.8) | 0.042 |
| Deployed with combat | 1049 | (11.5) | 207 | (13.1) | 842 | (11.2) | |
| Deployed without combat | 1339 | (14.7) | 212 | (13.4) | 1127 | (15.0) | |
| Positive screen for PTSD‡ | | | | | | | |
| No | 8414 | (92.5) | 1404 | (88.9) | 7010 | (93.2) | <0.001 |
| Yes | 685 | (7.5) | 175 | (11.1) | 510 | (6.8) | |
| Positive screen for depression§ | | | | | | | |
| No | 8578 | (94.3) | 1433 | (90.8) | 7145 | (95.0) | <0.001 |
| Yes | 521 | (5.8) | 146 | (9.3) | 375 | (5.0) | |
| Positive screen for panic or anxiety disorder¶ | | | | | | | |
| No | 8576 | (94.3) | 1426 | (90.3) | 7150 | (95.1) | <0.001 |
| Yes | 523 | (5.8) | 153 | (9.7) | 370 | (4.9) | |
| Occupational category (prior to separation) | | | | | | | |
| Combat specialist | 1129 | (12.4) | 215 | (13.6) | 914 | (12.2) | <0.001 |
| Officer or executive | 729 | (8.0) | 81 | (5.1) | 648 | (8.6) | |
| Healthcare | 885 | (9.7) | 183 | (11.6) | 702 | (9.3) | |
| Electronic equipment repairer or other technical | 2627 | (28.9) | 426 | (27.0) | 2201 | (29.3) | |
| Communications/intelligence | 860 | (9.5) | 131 | (8.3) | 729 | (9.7) | |
| Functional support and administration | 1734 | (19.1) | 328 | (20.8) | 1406 | (18.7) | |
| Craft worker | 229 | (2.5) | 36 | (2.3) | 193 | (2.6) | |
| Service and supply | 742 | (8.2) | 149 | (9.4) | 593 | (7.9) | |
| Student, trainee, or other | 164 | (1.8) | 30 | (1.9) | 134 | (1.8) | |
| Sex | | | | | | | |
| Male | 6826 | (75.0) | 1010 | (64.0) | 5816 | (77.3) | <0.001 |
| Female | 2273 | (25.0) | 569 | (36.0) | 1704 | (22.7) | |
| Period of study enrolment | | | | | | | |
| 2001–2003 (panel 1) | 7871 | (86.5) | 1245 | (78.9) | 6626 | (88.1) | <0.001 |
| 2004–2006 (panel 2) | 1228 | (13.5) | 334 | (21.2) | 894 | (11.9) | |
| Disabling illness or injury | | | | | | | |
| No | 8232 | (90.5) | 1356 | (85.9) | 6876 | (91.4) | <0.001 |
| Yes | 867 | (9.5) | 223 | (14.1) | 644 | (8.6) | |
| Years separated prior to reported employment status | | | | | | | |
| Less than 1 year | 2732 | (30.0) | 643 | (40.7) | 2089 | (27.8) | <0.001 |
| 1–2 years | 3296 | (36.2) | 527 | (33.4) | 2769 | (36.8) | |
| More than 2 years | 3071 | (33.8) | 409 | (25.9) | 2662 | (35.4) | |
| Age at military separation | | | | | | | |
| Mean (SD) | 37.3 | (9.1) | 36.1 | (10.4) | 37.5 | (8.8) | <0.001 |
| <25 years | 1203 | (13.2) | 316 | (20.0) | 887 | (11.8) | |
| 25–37 years | 2216 | (24.4) | 431 | (27.3) | 1785 | (23.7) | |
| 38–40 years | 1688 | (18.6) | 194 | (12.3) | 1494 | (19.9) | |
| 41–45 years | 2444 | (26.9) | 321 | (20.3) | 2123 | (28.2) | |
| >45 years | 1548 | (17.0) | 317 | (20.1) | 1231 | (16.4) | |
| Highest level of education achieved | | | | | | | |
| Less than bachelor's degree | 6842 | (75.2) | 1261 | (79.9) | 5581 | (74.2) | <0.001 |
| Bachelor's degree or higher | 2257 | (24.8) | 318 | (20.1) | 1939 | (25.8) | |
| Race/ethnicity | | | | | | | |
| Non-Hispanic white | 6136 | (67.4) | 1008 | (63.8) | 5128 | (68.2) | <0.001 |
| Non-Hispanic black | 1170 | (12.9) | 264 | (16.7) | 906 | (12.1) | |
| Other | 1793 | (19.7) | 307 | (19.4) | 1486 | (19.8) | |

Continued

Table 1 Continued

| Baseline* characteristics | Total separated N=9099 (100%) n (%)† | | Unemployed n=1579 (17%) n (%) | | Employed n=7520 (83%) n (%) | | p Value |
|-------------------------------------|---|--------|--|--------|--------------------------------------|--------|---------|
| Military pay grade | | | | | | | |
| Enlisted | 7133 | (78.4) | 1303 | (82.5) | 5830 | (77.5) | <0.001 |
| Officer | 1966 | (21.6) | 276 | (17.5) | 1690 | (22.5) | |
| Service branch | | | | | | | |
| Army | 3600 | (39.6) | 641 | (40.6) | 2959 | (39.4) | 0.278 |
| Navy/Coast Guard | 2379 | (26.2) | 425 | (26.9) | 1954 | (26.0) | |
| Air Force | 2419 | (26.6) | 407 | (25.8) | 2012 | (26.8) | |
| Marine Corps | 701 | (7.7) | 106 | (6.7) | 595 | (7.9) | |
| Marital status | | | | | | | |
| Never married | 2695 | (29.6) | 639 | (40.5) | 2056 | (27.3) | <0.001 |
| Married | 5964 | (65.6) | 858 | (54.3) | 5106 | (67.9) | |
| Divorced, separated, or widowed | 440 | (4.8) | 82 | (5.2) | 358 | (4.8) | |
| Smoking status | | | | | | | |
| Never smoker | 4918 | (54.1) | 829 | (52.5) | 4089 | (54.4) | <0.001 |
| Past smoker | 2484 | (27.3) | 389 | (24.6) | 2095 | (27.9) | |
| Current smoker | 1697 | (18.7) | 361 | (22.9) | 1336 | (17.8) | |
| Alcohol-related problems§ | | | | | | | |
| No | 8179 | (89.9) | 1407 | (89.1) | 6772 | (90.1) | 0.257 |
| Yes | 920 | (10.1) | 172 | (10.9) | 748 | (10.0) | |
| Life stressor score** | | | | | | | |
| Low/mild stress | 8211 | (90.2) | 1347 | (85.3) | 6864 | (91.3) | <0.001 |
| Moderate stress | 743 | (8.2) | 193 | (12.2) | 550 | (7.3) | |
| Major stress | 145 | (1.6) | 39 | (2.5) | 106 | (1.4) | |
| PCS score¶ | | | | | | | |
| <15th percentile (poor health) | 2180 | (24.0) | 514 | (32.6) | 1666 | (22.2) | <0.001 |
| 15th–85th percentile (average) | 5978 | (65.7) | 896 | (56.7) | 5082 | (67.6) | |
| >85th percentile (excellent health) | 941 | (10.3) | 169 | (10.7) | 772 | (10.3) | |

*Baseline characteristics were assessed using the most recent survey submitted prior to separation from service.

†Percents may not sum to 100% due to rounding.

‡Evaluated using the 17-item PTSD Checklist-Civilian Version.

§Assessed using the PRIME-MD Patient Health Questionnaire.

¶Measurement of physical health based on the Medical Outcomes Study Short Form 36-Item Health Survey for Veterans.

**Measurement based on the Holmes and Rahe Social Readjustment Rating Scale.

PCS, physical component summary; PTSD, post-traumatic stress disorder.

with PTSD symptoms was twice as high compared to those without PTSD (29.4% vs. 15.4%). This finding suggests that PTSD was likely more of an impediment to finding a job than time for this particular group.

Former Navy and Air Force personnel who routinely retired, had higher odds of unemployment compared with the Army. When we examined other characteristics of former Navy and Air Force members, we found few differences other than proportionally more Navy and Air Force veterans in this stratum retired at a lower pay grade than Army veterans, refuting the potential hypothesis that these service members retired with a more adequate pension for ordinary living. The reason for this finding is unknown; thus, more research is needed to understand service-specific differences in employment paths of retired veterans, which may be possible using data from future Millennium Cohort surveys which will be collecting information on job seeking.

Two important groups, women and non-Hispanic black participants, were at increased odds of unemployment following separation from service. Similarly, the BLS reported that African-Americans had a higher unemployment rate (14.7%) compared with Caucasians (10.7%).³¹ By contrast, the March 2011 BLS news release on the employment situation of veterans found

little difference in the unemployment rate of male and female service members who joined the military after September 2001 (11.4% and 12.0%, respectively).³¹ Further study is warranted to determine why women and black veterans in some cohorts have higher unemployment, when all, theoretically, had equal access to training and opportunity for advancement within the military system.

Current smoking was associated with not having a job among the routinely retired, and no significant association was found between alcohol abuse and job status in multivariable or univariate analyses. Former service members who separated for other reasons and reported alcohol-related problems prior to separation had marginally significantly lower odds of reporting unemployment at follow-up. Though insignificant, this finding appears surprising, but may be explained by working individuals who may misuse alcohol to cope with the stress of their daily responsibilities. It is also important to note that while alcohol-related problems are reported, these are not a necessary indication of functional impairment.

The proportion unemployed in this population is concerning; yet it should not be compared with the national rate, which includes those actively seeking employment as its denominator,³⁷ and which we could not determine for our

Workplace

Table 2 Multivariable model of the adjusted odds of unemployment post service by reason for military separation among Millennium Cohort participants (N=9099)

| Baseline† characteristics | Reason for military separation | | | | | |
|---|--------------------------------|----------------|------------------------------------|----------------|------------------------|----------------|
| | Routine retirement (N=5488) | | Fulfilled service term (N=2165) | | Other reasons (N=1446) | |
| | OR | (95% CI) | OR | (95% CI) | OR | (95% CI) |
| Deployment experience | | | | | | |
| Not recently deployed | 1.00 | | 1.00 | | 1.00 | |
| Deployed with combat | 1.14 | (0.87 to 1.50) | 1.20 | (0.87 to 1.64) | 1.00 | (0.67 to 1.48) |
| Deployed without combat | 0.86 | (0.68 to 1.08) | 1.20 | (0.88 to 1.64) | 1.09 | (0.70 to 1.71) |
| Positive screen for PTSD‡ | | | | | | |
| No | 1.00 | | 1.00 | | 1.00 | |
| Yes | 0.71 | (0.46 to 1.09) | 1.25 | (0.82 to 1.91) | 1.08 | (0.72 to 1.62) |
| Positive screen for depression§ | | | | | | |
| No | 1.00 | | 1.00 | | 1.00 | |
| Yes | 1.67* | (1.05 to 2.63) | 0.87 | (0.53 to 1.44) | 1.02 | (0.65 to 1.60) |
| Positive screen for panic or anxiety disorder§ | | | | | | |
| No | 1.00 | | 1.00 | | 1.00 | |
| Yes | 1.63* | (1.10 to 2.43) | 0.97 | (0.58 to 1.62) | 1.45 | (0.93 to 2.26) |
| Occupation category (prior to separation) | | | | | | |
| Functional support and administration | 1.00 | | 1.00 | | 1.00 | |
| Combat specialist | 1.25 | (0.91 to 1.71) | 0.84 | (0.56 to 1.25) | 1.14 | (0.72 to 1.80) |
| Officer or executive | 0.92 | (0.65 to 1.32) | 0.82 | (0.29 to 2.33) | 0.44 | (0.15 to 1.25) |
| Healthcare | 1.27 | (0.95 to 1.70) | 0.69 | (0.43 to 1.10) | 0.72 | (0.44 to 1.17) |
| Electronic equipment repairer or other technical | 1.03 | (0.82 to 1.31) | 0.68* | (0.47 to 0.98) | 0.71 | (0.47 to 1.05) |
| Communications/intelligence | 0.88 | (0.62 to 1.25) | 0.64* | (0.41 to 0.99) | 0.75 | (0.45 to 1.25) |
| Craft worker | 1.01 | (0.60 to 1.72) | 0.65 | (0.28 to 1.52) | 0.56 | (0.24 to 1.33) |
| Service and supply | 1.21 | (0.87 to 1.69) | 0.92 | (0.58 to 1.46) | 0.79 | (0.49 to 1.27) |
| Student, trainee, or other | 1.67 | (0.73 to 3.79) | 0.71 | (0.36 to 1.42) | 0.93 | (0.40 to 2.20) |
| Sex | | | | | | |
| Male | 1.00 | | 1.00 | | 1.00 | |
| Female | 1.70* | (1.38 to 2.10) | 1.42* | (1.09 to 1.84) | 1.17 | (0.88 to 1.56) |
| Period of study enrolment | | | | | | |
| 2001–2003 (panel 1) | N/A¶ | | 1.00 | | 1.00 | |
| 2004–2006 (panel 2) | N/A¶ | | 1.33* | (1.03 to 1.70) | 1.63* | (1.21 to 2.18) |
| Disabling illness or injury | | | | | | |
| No | 1.00 | | 1.00 | | 1.00 | |
| Yes | 1.59* | (1.20 to 2.10) | 1.60* | (1.06 to 2.41) | 0.92 | (0.63 to 1.33) |
| Years separated prior to reported employment status | | | | | | |
| Less than 1 year | 1.00 | | 1.00 | | 1.00 | |
| 1–2 years | 0.57* | (0.47 to 0.68) | 0.50* | (0.38 to 0.65) | 0.52* | (0.38 to 0.71) |
| More than 2 years | 0.43* | (0.35 to 0.53) | 0.39* | (0.29 to 0.52) | 0.38* | (0.27 to 0.52) |
| Age at military separation (10-year intervals) | 2.53* | (2.08 to 3.07) | 0.76 | (0.56 to 1.03) | 1.39* | (1.07 to 1.82) |
| Highest level of education achieved | | | | | | |
| Less than bachelor's degree | 1.00 | | 1.00 | | 1.00 | |
| Bachelor's degree or higher | 0.70* | (0.53 to 0.93) | 0.77 | (0.41 to 1.44) | 0.48* | (0.25 to 0.90) |
| Race/ethnicity | | | | | | |
| Non-Hispanic white | 1.00 | | 1.00 | | 1.00 | |
| Non-Hispanic black | 1.31* | (1.04 to 1.65) | 1.82* | (1.27 to 2.63) | 1.55* | (1.06 to 2.28) |
| Other | 1.14 | (0.89 to 1.45) | 1.13 | (0.84 to 1.52) | 1.38 | (0.98 to 1.96) |
| Military pay grade | | | | | | |
| Enlisted | 1.00 | | 1.00 | | 1.00 | |
| Officer | 1.10 | (0.81 to 1.50) | 0.54 | (0.23 to 1.29) | 0.90 | (0.40 to 2.02) |
| Service branch | | | | | | |
| Army | 1.00 | | 1.00 | | 1.00 | |
| Navy/Coast Guard | 1.38* | (1.10 to 1.75) | 1.13 | (0.83 to 1.54) | 0.76 | (0.54 to 1.07) |
| Air Force | 1.46* | (1.16 to 1.83) | 1.24 | (0.89 to 1.73) | 1.29 | (0.91 to 1.83) |
| Marine Corps | 0.91 | (0.57 to 1.44) | 0.84 | (0.59 to 1.19) | 1.08 | (0.61 to 1.90) |

Continued

Table 2 Continued

| Baseline† characteristics | Reason for military separation | | | | | |
|-------------------------------------|--------------------------------|----------------|------------------------------------|----------------|------------------------|----------------|
| | Routine retirement (N=5488) | | Fulfilled service term (N=2165) | | Other reasons (N=1446) | |
| | OR | (95% CI) | OR | (95% CI) | OR | (95% CI) |
| Marital status | | | | | | |
| Never married | 1.00 | | 1.00 | | 1.00 | |
| Married | 0.55* | (0.43 to 0.71) | 0.68* | (0.51 to 0.90) | 0.91 | (0.68 to 1.22) |
| Divorced, separated, or widowed | 0.63* | (0.43 to 0.93) | 0.55 | (0.24 to 1.24) | 0.73 | (0.35 to 1.52) |
| Smoking status | | | | | | |
| Never smoker | 1.00 | | 1.00 | | 1.00 | |
| Past smoker | 0.91 | (0.76 to 1.10) | 0.88 | (0.65 to 1.18) | 1.15 | (0.83 to 1.61) |
| Current smoker | 1.40* | (1.11 to 1.76) | 0.93 | (0.70 to 1.24) | 1.36 | (1.00 to 1.87) |
| Alcohol-related problems§ | | | | | | |
| No | 1.00 | | 1.00 | | 1.00 | |
| Yes | 1.28 | (0.91 to 1.81) | 0.87 | (0.64 to 1.19) | 0.70 | (0.48 to 1.00) |
| Life stressor score** | | | | | | |
| Low/mild stress | 1.00 | | 1.00 | | 1.00 | |
| Moderate stress | 1.23 | (0.92 to 1.65) | 1.80* | (1.22 to 2.66) | 1.17 | (0.81 to 1.70) |
| Major stress | 0.92 | (0.48 to 1.75) | 1.11 | (0.39 to 3.13) | 1.17 | (0.59 to 2.34) |
| PCS score†† | | | | | | |
| <15th percentile (poor health) | 1.57* | (1.30 to 1.89) | 1.06 | (0.77 to 1.45) | 1.64* | (1.23 to 2.19) |
| 15th–85th percentile (average) | 1.00 | | 1.00 | | 1.00 | |
| >85th percentile (excellent health) | 1.17 | (0.87 to 1.56) | 1.11 | (0.81 to 1.53) | 1.14 | (0.74 to 1.77) |

*Indicates significance at the $p < 0.05$ level.

†Baseline characteristics were assessed using the most recent survey submitted prior to separation from service.

‡Evaluated using the 17-item PTSD Checklist-Civilian Version.

§Assessed using the PRIME-MD Patient Health Questionnaire.

¶This variable was excluded from this model due to insufficient numbers in panel 2, because these participants are generally younger and ineligible for retirement.

**Measurement based on the Holmes and Rahe Social Readjustment Rating Scale.

††Measurement of physical health based on the Medical Outcomes Study Short Form 36-Item Health Survey for Veterans.

N/A, not applicable; PCS, physical component summary; PTSD, post-traumatic stress disorder.

study population. The routinely retired group had the lowest proportion without a job. The mean age of this group was 43 years at the time of separation, and all but two were below the age of 65 years. Thus, these individuals may still desire to work after retiring from the military. The valuable work experience, resources and skills gained during 20 or more years in the military may make these individuals more attractive to employers than others. On the other hand, those separated for other reasons may be the least attractive to employers, or they may not be seeking work, or they may be physically incapable of working. Many were receiving severance pay, as we noted that nearly 50% of this group separated because of a disability and service-connected disabilities with a rating >30% entitle service members who have served less than 20 years to retirement benefits.³⁸ Additional analyses confirmed that separating from service due to a disability was univariately associated with unemployment postservice (OR, 1.60; 95%CI, 1.26–2.03) (data not shown). Reporting a prior disability did not significantly affect odds of unemployment among this group; however, this may have been due to over-adjustment issues or the diversity of this “other reasons” group. Also, given the high percentage of disability among this group, we linked participant records with the Joint Theater Trauma Registry to examine deployment-related injuries prior to separation. Few injuries were recorded, and no significant associations between deployment injuries and employment status were found in additional analyses; however, lack of significance may have been due to reduced

power. Since disabling illnesses or injuries reported at baseline and poor physical health significantly affected unemployment in two-thirds of our groups, further studies are needed to evaluate the impact of military-related injuries on postservice employability.

There were some limitations to this study that should be considered. We could not determine whether the unemployed had been unsuccessfully searching for employment, had recently lost a job, had trouble keeping jobs, were in school/training, were parenting, or were physically unable to work; although separation codes provided some insight. Those who separated for other reasons, in particular, represented a more diverse group of former service members than those who routinely retired or finished their service term. In addition, the number of civilian jobs held or lost since the date of separation was not captured. Personnel files do not provide a detailed description for the reason for separation, so some individuals may have been misclassified based on inaccuracies in these data, or slight differences across service branch separation policies. Survey non-response due to attrition may have also been an issue in this population of separated personnel. Previous research has shown little follow-up non-response bias in the cohort as a whole; however, personnel who separated for reasons other than retirement had a higher probability of follow-up non-response.³⁹ Also, health outcomes were self-reported, but previous research on this cohort has shown participant data to be reliable.^{13 25 40} Mental disorders and behaviours were measured prior to military separation as predisposing conditions;

Workplace

however, they may fluctuate in severity over time. Whether they would have limited the ability to obtain or hold employment at the time of separation was not determined in this survey. PTSD, depression and panic/anxiety were identified based on self-reported screening tools, not clinical diagnoses. However, standardised instruments with high sensitivity and specificity were used to reduce misclassification error. Although this study population may differ from the full population of recent veterans, research has shown the cohort to be well representative of the US military.¹²

The Millennium Cohort Study provided a unique opportunity to investigate a robust sample of former military personnel and make inferences across all service branches, increasing the generalisability of the findings. The longitudinal design allowed for prospective follow-up of postservice employment, while adjusting for many characteristics. Finally, stigma attached with mental disorders may prevent some individuals from seeking care within the Military Health System; therefore, use of confidential surveys eliciting self-reported symptoms may have allowed for greater capture of mental health morbidity.

CONCLUSION

Employment is influenced by mental and physical health, and may be considered a strong marker of overall health. Not surprisingly, the reason for separation from military service is a strong indicator of postservice employment status. The degree to which reasons for separation vary make investigation of this population complex. Some wish to retire; others seek military occupations for the promise of furthering their education post-service, or for superior on-the-job training, to position them well for future employment. Previous research highlighted retention issues surrounding mental disorders and suggested a higher exit prevalence from the military among those with mental health needs.³ The current study suggests that screening positive for certain mental disorders like panic/anxiety or depression are associated with reporting unemployment among the routinely retired group, but being deployed during service was not associated with unemployment in any of the groups examined. These data offer a first look at subgroups that may be disadvantaged in achieving postservice employment, though the preponderance of information suggests a military workforce able to find employment after serving. Continued research is necessary to investigate other impactful military-related experiences not measured in this study, and to understand how in-service training and education in subpopulations will influence employment in the future.

Acknowledgements In addition to the authors, the Millennium Cohort Study Team includes Melissa Bagnell, Gina Creaven, Nancy Crum-Cianflone, James Davies, Nisara Granado, Gia Gumbs, Lesley Henry, Dennis Hernando, Andrea Ippolito, Kelly Jones, Lauren Kipp, Cynthia LeardMann, William Lee, Michelle Linfesty, Gordon Lynch, Hope McMaster, Sheila Medina-Torne, Amanda Pietrucha, Teresa Powell, Kari Sausedo, Amber Seelig, Beverly Sheppard, Katherine Snell, Steven Speigle, Jennifer Walstrom, John Wesner, Martin White and James Whitmer. We are indebted to the Millennium Cohort Study participants, without whom these analyses would not be possible. We appreciate the support from the Management Information Division, US Defense Manpower Data Center (Seaside, CA, USA); Military Operational Medicine Research Program and US Army Medical Research and Materiel Command (Fort Detrick, MD, USA). In addition, we thank Michelle LeWark from the Naval Health Research Center (San Diego, CA, USA).

Funding The Millennium Cohort Study is funded through the Military Operational Medicine Research Program, USA Army Medical Research and Materiel Command (Fort Detrick, MD, USA). Resources from the VA Puget Sound Health Care System supported Dr Boyko's involvement in this research. The funding organisation had no role in the design and conduct of the study; collection, analysis, or preparation of data; or preparation, review or approval of the manuscript.

Disclaimer This represents report 11–49 supported by the Department of Defense, under work unit number 60002. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of Defense, or the US Government. Approved for public release; distribution is unlimited. This research was conducted in compliance with all applicable federal regulations governing the protection of human subjects (protocol NHRC.2000.0007).

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Britt TW, Greene-Shortridge TM, Castro CA. The stigma of mental health problems in the military. *Mil Med* 2007;172:157–61.
- 2 Strayer R, Ellenhorn L. Vietnam Veterans—Study Exploring Adjustment Patterns and Attitudes. *J Soc Issues* 1975;31:81–93.
- 3 Hoge CW, Auchterlonie JL, Milliken CS. Mental health problems, use of mental health services, and attrition from military service after returning from deployment to Iraq or Afghanistan. *JAMA* 2006;295:1023–32.
- 4 Hoge CW, Castro CA, Messer SC, et al. Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *N Engl J Med* 2004;351:13–22.
- 5 Milliken CS, Auchterlonie JL, Hoge CW. Longitudinal assessment of mental health problems among active and reserve component soldiers returning from the Iraq War. *JAMA* 2007;298:2141–8.
- 6 Smith TC, Ryan MA, Wingard DL, et al. New onset and persistent symptoms of post-traumatic stress disorder self reported after deployment and combat exposures: prospective population based US military cohort study. *BMJ* 2008;336:366–71.
- 7 *Healthcare Inspection: review of the veterans health administration's use of appropriated funds for research*. Washington, DC: Department of Veterans Affairs, Office of Inspector General; 2009. Report No.: 08-03007-80.
- 8 Anthony WA, Rogers ES, Cohen M, et al. Relationships between psychiatric symptomatology, work skills, and future vocational performance. *Psychiatr Serv* 1995;46:353–8.
- 9 Crowther RE, Marshall M, Bond GR, et al. Helping people with severe mental illness to obtain work: systematic review. *BMJ* 2001;322:204–8.
- 10 Gray GC, Chesbrough KB, Ryan MAK, et al. The Millennium Cohort Study: a 21-year prospective cohort study of 140,000 military personnel. *Mil Med* 2002;167:483–8.
- 11 Riddle JR, Smith TC, Smith B, et al. Millennium Cohort: the 2001–2003 baseline prevalence of mental disorders in the U.S. military. *J Clin Epidemiol* 2007;60:192–201.
- 12 Ryan MA, Smith TC, Smith B, et al. Millennium Cohort: enrollment begins a 21-year contribution to understanding the impact of military service. *J Clin Epidemiol* 2007;60:181–91.
- 13 Smith B, Wingard DL, Ryan MA, et al. U.S. military deployment during 2001–2006: comparison of subjective and objective data sources in a large prospective health study. *Ann Epidemiol* 2007;17:976–82.
- 14 Burnett-Zeigler I, Valenstein M, Ilgen M, et al. Civilian Employment Among Recently Returning Afghanistan and Iraq National Guard Veterans. *Mil Med* 2011;176:639–46.
- 15 Cohen S, Suri P, Amick M, et al. Clinical and demographic factors associated with employment status in US military veterans returning from Iraq and Afghanistan. *Work* 2013;44(2):213–19.
- 16 Dillman DA, Phelps G, Tortora R, et al. Response rate and measurement differences in mixed-mode surveys using mail, telephone, interactive voice response (IVR) and the Internet. *Soc Sci Res* 2009;38:3–20.
- 17 *Military attrition: better data, coupled with policy changes, could help the services reduce early separations*. Washington, DC: US General Accounting Office Report; July 2000. Report No.: GAO/NSIAD-98-213.
- 18 Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ Primary Care Study. Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire. *JAMA* 1999;282:1737–44.
- 19 Spitzer RL, Williams JB, Kroenke K, et al. Utility of a new procedure for diagnosing mental disorders in primary care. The PRIME-MD 1000 Study. *JAMA* 1994;272:1749–56.
- 20 Holmes TH, Rahe RH. Social Readjustment Rating Scale. *J Psychosom Res* 1967;11:213.
- 21 Ware JE, Sherbourne CD. The Mos 36-Item Short-Form Health Survey(Sf-36).1. Conceptual-framework and item selection. *Med Care* 1992;30:473–83.
- 22 Weathers FW, Litz BT, Herman DS, et al. eds. The PTSD Checklist (PCL): reliability, validity, and diagnostic utility. Paper presented at the Annual Meeting of International Society for Traumatic Stress Studies; October 1993, San Antonio, Texas.

- 23 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th edn. DSM-IV. Washington, DC: American Psychiatric Association, 1994.
- 24 Brewin CR. Systematic review of screening instruments for adults at risk of PTSD. *J Trauma Stress* 2005;18:53–62.
- 25 Smith TC, Smith B, Jacobson IG, *et al*. Reliability of standard health assessment instruments in a large, population-based cohort study. *Ann Epidemiol* 2007;17:271–84.
- 26 Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med* 2001;16:606–13.
- 27 Fann JR, Bombardier CH, Dikmen S, *et al*. Validity of the Patient Health Questionnaire-9 in assessing depression following traumatic brain injury. *J Head Trauma Rehabil* 2005;20:501–11.
- 28 Glantz S, Slinker B. *Primer of applied regression and analysis of variance*. New York, NY: McGraw-Hill, 1990.
- 29 Wells TS, LeardMann CA, Fortuna SO, *et al*. A Prospective Study of Depression Following Combat Deployment in Support of the Wars in Iraq and Afghanistan. *Am J Public Health* 2010;100:90–9.
- 30 Clark J. Subcommittee on Economic Opportunity holds Hearing on Veterans Employment and Training Service Budget Request. 2011. <http://veterans.house.gov/news/PRArticle.aspx?NewsID=2294> (accessed 5 Apr 2011).
- 31 BLS. Employment situation of veterans—2010. Dept of Labor; 11 March 2011.
- 32 Lauber C, Bowen JL. Low mood and employment: When affective disorders are intertwined with the workplace—A UK perspective. *Int Rev Psychiatry* 2010;22:173–82.
- 33 Lerner D, Henke RM. What does research tell us about depression, job performance, and work productivity? *J Occup Environ Med* 2008;50:401–10.
- 34 Uutela A. Economic crisis and mental health. *Curr Opin Psychiatry* 2010;23:127–30.
- 35 Resnick SG, Rosenheck RA. Posttraumatic stress disorder and employment in veterans participating in Veterans Health Administration Compensated Work Therapy. *J Rehabil Res Dev* 2008;45:427–35.
- 36 Geuze E, Vermetten E, de Kloet CS, *et al*. Neuropsychological performance is related to current social and occupational functioning in veterans with posttraumatic stress disorder. *Depress Anxiety* 2009;26:7–15.
- 37 How the government measures unemployment. U.S. Bureau of Labor Statistics Division of Labor Force Statistics. 2009. http://www.bls.gov/cps/cps_htgm.htm#unemployed
- 38 Army Regulation 635-40: Physical Evaluation for Retention, Retirement, or Separation. Washington, DC: US Department of the Army, Aug 15, 1990.
- 39 Littman AJ, Boyko EJ, Jacobson IG, *et al*. Assessing nonresponse bias at follow-up in a large prospective cohort of relatively young and mobile military service members. *Bmc Med Res Methodol* 2010;10.
- 40 Smith TC, Jacobson IG, Smith B, *et al*. The occupational role of women in military service: validation of occupation and prevalence of exposures in the Millennium Cohort Study. *Int J Environ Health Res* 2007;17:271–84.



The impact of prior deployment experience on civilian employment after military service

Jaime L Horton, Isabel G Jacobson, Charlene A Wong, Timothy S Wells, Edward J Boyko, Besa Smith, Margaret A K Ryan, Tyler C Smith and for the Millennium Cohort Study Team

Occup Environ Med 2013 70: 408-417 originally published online March 21, 2013

doi: 10.1136/oemed-2012-101073

Updated information and services can be found at:
<http://oem.bmj.com/content/70/6/408>

These include:

References

This article cites 30 articles, 2 of which you can access for free at:
<http://oem.bmj.com/content/70/6/408#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>

REPORT DOCUMENTATION PAGE

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB Control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

| | | | | | |
|--|----------------------------|--|---|--|--|
| 1. REPORT DATE (DD MM YY) 09 09 11 | | 2. REPORT TYPE Journal Article | | 3. DATES COVERED (from – to) 2001–2008 | |
| 4. TITLE Civilian Employment After Military Service | | | | 5a. Contract Number: 5b. Grant Number: 5c. Program Element Number: 5d. Project Number: 5e. Task Number: 5f. Work Unit Number: 60002 | |
| 6. AUTHORS Horton, Jaime L.; Isabel G. Jacobson, Charlene A. Wong, Timothy S. Wells, Edward J. Boyko, Besa Smith, Margaret A.K. Ryan, & Tyler C. Smith for the Millennium Cohort Study Team | | | | | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Commanding Officer Naval Health Research Center 140 Sylvester Rd San Diego, CA 92106-3521 | | | | | |
| 8. SPONSORING/MONITORING AGENCY NAMES(S) AND ADDRESS(ES) Commanding Officer Naval Medical Research Center 503 Robert Grant Ave Silver Spring, MD 20910-7500 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER Report No. 11-49 | |
| | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) NMRC/BUMED | |
| Chief, Bureau of Medicine and Surgery (MED 00), Navy Dept 2300 E Street NW Washington, DC 20372-5300 | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(s) | |
| | | | | | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited. | | | | | |
| 13. SUPPLEMENTARY NOTES <u>Occupational and Environmental Medicine</u> (2013), <u>70</u> , 408–417 | | | | | |
| 14. ABSTRACT <p>Concerns exist over whether mental disorders and combat deployment experience among US military personnel affect postservice employment. The objective was to determine if deployment experience in support of the operations in Iraq and Afghanistan or a positive screen for posttraumatic stress disorder (PTSD), depression, or panic/anxiety disorders were associated with reporting employment after military service. Former US military active duty and Reserve or National Guard personnel from the Millennium Cohort Study were followed for an average of 3.2 years from the time of mental disorder reporting and potential deployment experience to self-reported employment status.</p> <p>Among all separated personnel except for those who retired with a pension, screening positive for depression, panic, or anxiety did not significantly affect the ability to obtain future civilian employment. Military combat experiences examined in this study and a positive screen for PTSD were not associated with postservice employment status. Screening positive for depression and panic/anxiety disorders predicted higher odds of subsequent unemployment among those who routinely retired with a pension. Most veterans secured civilian careers in fields similar to their previous military occupations.</p> | | | | | |
| 15. SUBJECT TERMS employment, military personnel | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT UNCL | 18. NUMBER OF PAGES 11 | 18a. NAME OF RESPONSIBLE PERSON Commanding Officer |
| a. REPORT UNCL | b. ABSTRACT UNCL | c. THIS PAGE UNCL | | | 18b. TELEPHONE NUMBER (INCLUDING AREA CODE) COMM/DSN: (619) 553-8429 |